An Inaugural
Spay Popul april
19 1824
10 9 18 14
Cholera Morbus

By

George Williamson Richmond Ninginia


Copioces discharges of bilious matter up wards and arounwards, accompanied with gripings and Spason, constitute Cholora Mor. bus. — This disease is placed by Cellow in the class necesses and order Spasoni, and by Doctor Chapman it is arranged among the diseases of the digestive System.

The cause of this disease are the same as those which produce its kindred affections. Willows colored by marsh mian made, in other townsomers, suddenly succeeded by colo, intemperance in eating and drinking be; but it is more than probable that these last only act as existing causes on a system almady strongly predisposed to an attack of the disease from the provinces on morha action produced by tharsh the

Like other gastric affections, this disease

comes on with nausia, painful sensations about the epigastric region, followed by vomiting and purging, first of the contents of the Stomach and bowels and afterwards of buro Bile, attended with griping pains and sparmodic affections of the muscles of the abde men: the pulso is weak and fluttering and the strengthe considerably prostrated: as the disease advances, all the above symtoms be come much aggravated, the debitily becomes extreme, the vomitting and purging unremitting, the pulse weak and almost imperceptible, the surface cold and palled, the respiration hurried, with great anxiety and restlepness, attended by cramps of the muscles of the extrem ities, and should the patient not be very soon releaved, death soon puts an end to the appole

The Diagnoses of Cholera are plain: it is distinguished from depending and diarrham by



The executions pensisting of pero bile unmixed with blood and facularly matter and from Colic by, the absence of conslipation.

The Brognosis is unfavorable when there are faintings, edit sweats, great at belity phosecoughs and Spasons: the absence of the unfavorable symtoms of course shew the absence of immediate danger.

Of the Fathology of this desease I then to there can be bet little doubt of its being, a gastric affection and extensive sympathelically to the stover inclined to concern with Doctor Chapman as lotte primary seat, from the following considerations org: the stomach show the first symbons of an attack, the first discharges are not bete, but consist simply of the contents of the stomach and not litt writation of the stomachias lasted sometime, do the believes executations appear. There are frequently collicious executations appear. There are frequently collicious executations appear without for a decing Choleran Mithough judging from appear.

case the bele is much more vitiated in this discase than usual, I think in fact we should consider it more bland and in offensive than in the ordenary states considering the vapidity, with which it is secretic and how soon it paper off

Believing Doctor Chapman's theory of the disease to be cornet I shall follow kim in the Freatment.

At the commencement of the dis case opium will in many cases completely by arrest and overcome way bad symbols get in most cases I thenk it advisable to rid the stomach of its initiating continuts by an emitic of Spicaceanha followed by copia our draughts of spicaceanha followed by copia our draughts of spicaceanha followed by copia our draughts of spicaceanha debility be fore that I should move of treating the drawes by copious draughts of warm flueds was lame pering with a sident distance with feells and inefficient veneralises; and as their object was

to empty the stomach I can not conceive why they requested he more cortain and active measures to protecting the effect. I think the Succeeding the in this case is betiefted of the openior not only of an oblying the stomach; but I or leving the Spanns and promoting a determination to the skin, which we desirable objects.

If the pain be violent and the patient with a war it, remiscolions house not be resorted to; but we can not be governed on this as in most disease by the pulser as to the hordisety or the extent to which it should be carried. The general aspired of the case will common by indicate or forted the west of the tancel. In doubtful cases it would be better to detend a small grand by and watch the iffice I, and if the pulser mon from it, which is not uncommon, we can repeat at pleasure. The warm hath is very of pleastle to this disease, more especially, when the lancet is



indicated by the pain but can not be used on account of debetity. It helps to releve spasm and produce a general instant of a local excitement, it brings on an open form of the disease and thereby enables is to attack it without heitaling as to the property of our other remedies, more parlice early remisection.

The preceding owned is failing it will in proper, again to aught to opening given both by the mouth and as an Enema, or it may be given in lineture with the effect escal of aught and milk, an influeion of server water and milk, an influeion of serventaria, strong coffee without reasons sugar, example year their properties of hel water labelling reguently, should be touch succeptively, if he worniting continue obstinate. Calomet in small are repealed does will preguently second in allaying the writability of the stomach and



Broweld External applications to the region of the Stomach are heighly valuable and among the most so may be en une rated the Clove Bag, Mannels uningout of hot spirits of Surpentine, camphorated Spirits or strongly impregnated with the warmer spices, as perform, mustared to be about all own chief confidence who weld be reposed in Blitters placed over the stomach and applied to the internative sales if the general systems hould require it. If the disease advances to rapially for the drawing of a blister we may use the following.

The Nitrie Acid tone part

Mater. . . two parts \_ Alinen rag may be dipped in this and applied over he stomach for a moment or two: if it be loo wident in its action it may be mentralized by a colition of the Cart of Totask. Maving by these means checked the property

of the disease our attention should ment be directed to prevent a recurrence, to which the patient is very liable: to effect which weshould direct our patient to avoid most sedulous by the exciting causes, as exposure to the hot sun, night air, excepts in eating or drenking, violent exercise, and cold extremilies should be particularly quarded against. The wear ing of flannel next to the skin is of infinite in. portance and should not be overlooked-The Bowels should be kept in a soluble state by small doses of magnisia or while bast and tone imparted to the alimentary Canal by tonics; the best of which are the Quapia, Colombo and Elizor Vitriol - A diet light and mutritive should be selected -